

Unless otherwise specified, V is a vector space over a field \mathbb{F} .

#1 More Eigenfun Let $A, B, C, D, P \in \mathfrak{gl}_n(\mathbb{F})$. Suppose that D is a diagonal matrix with diagonal entries $\lambda_1, \dots, \lambda_n$ and that A is diagonalizable with eigenvalues $\lambda_1, \dots, \lambda_n$. In addition, suppose that P is invertible.

- Show that E_{ij} is an eigenvector for ad_D for each $i, j = 1, \dots, n$. What are ad_D 's eigenvalues? Why is ad_D diagonalizable?
- Let $A = PDP^{-1}$ and $B = PCP^{-1}$. Show that $[A, B] = P[D, C]P^{-1}$.
- (Grad Students)** Show that ad_A is diagonalizable.

#2 Direct Sums Recall the definition of a vector space direct sum: $L = A \oplus B$ for some subspaces A and B if $L = A + B$ (i.e. for every $\mathbf{v} \in L$ there exists some $\mathbf{a} \in A$ and $\mathbf{b} \in B$ such that $\mathbf{v} = \mathbf{a} + \mathbf{b}$) and $A \cap B = \{\mathbf{0}\}$.

If in addition L is a Lie algebra, A and B are subalgebras, and $[A, B] = \{\mathbf{0}\}$, then we say $L = A \oplus B$ as Lie algebras (i.e. L is a direct sum of its subalgebras A and B).

Let \mathbb{F} be field of characteristic 0. Show that $\mathfrak{gl}_n(\mathbb{F}) = \mathfrak{sl}_n(\mathbb{F}) \oplus \mathbb{F}I_n$ as Lie algebras (note: $\mathbb{F}I_n$ is the set of scalar multiples of the identity matrix).

#3 An Ideal Problem Let $X \subseteq L$ (X is just a subset). Recall that $C_L(X) = \{\mathbf{v} \in L \mid [\mathbf{v}, \mathbf{x}] = \mathbf{0} \text{ for all } \mathbf{x} \in X\}$ is the centralizer of X in L .

- Show that $C_L(X)$ is a subalgebra of L (don't forget to show it is a subspace as well).
- Assume $X \triangleleft L$. Show that $C_L(X) \triangleleft L$.

#4 Automorphism via Conjugation For some fixed $J \in \mathfrak{gl}_n(\mathbb{F})$, recall that $\mathfrak{gl}_n^J(\mathbb{F}) = \{X \in \mathfrak{gl}_n(\mathbb{F}) \mid JX + X^T J = 0\}$. Suppose that A is an invertible $n \times n$ matrix such that $A^T J A = J$. Show that $\varphi : \mathfrak{gl}_n^J(\mathbb{F}) \rightarrow \mathfrak{gl}_n^J(\mathbb{F})$ defined by $\varphi(X) = A^{-1} X A$ is an automorphism of $\mathfrak{gl}_n^J(\mathbb{F})$.

Note: The first thing you should verify is that φ actually maps from $\mathfrak{gl}_n^J(\mathbb{F})$ to itself.

#5 Derivations Recall $\text{Der}(\mathbb{F}[x]) = \{\partial : \mathbb{F}[x] \rightarrow \mathbb{F}[x] \mid \partial \text{ is linear and } \partial(fg) = \partial(f)g + f\partial(g) \text{ for all } f, g \in \mathbb{F}[x]\}$ where $\mathbb{F}[x]$ is the algebra of polynomials with coefficients in $\mathbb{F}[x]$.

- Let $\partial \in \text{Der}(\mathbb{F}[x])$. Show that $\partial(1) = 0$ and thus any derivation applied to a constant yields 0.
- Let $p(x) \in \mathbb{F}[x]$. Define $\partial(f(x)) = p(x)f'(x)$ (i.e. $\partial = p(x)\frac{d}{dx}$). Show that ∂ is a derivation on $\mathbb{F}[x]$.
- Compute $[p(x)\frac{d}{dx}, q(x)\frac{d}{dx}]$ where $p(x), q(x) \in \mathbb{F}[x]$.
- (Grad Students)** Show that $\text{Der}(\mathbb{F}[x]) = \{p(x)\frac{d}{dx} \mid p(x) \in \mathbb{F}[x]\}$ (i.e. all derivations on $\mathbb{F}[x]$ are like those defined in part (b)).