
- #1 Basics More "Basic" Calculations
  - (a) Compute  $e^z$  for  $z = -\frac{i\pi}{3}, \frac{1}{2} \frac{i\pi}{4}$ , and  $-1 + \frac{i3\pi}{2}$ .
  - (b) Compute Log(4+4i) and Log(-10i). Then also compute log(4+4i) and log(-10i).
  - (c) Compute  $(-1)^i$  and indicate what the principal branch yields.
  - (d) [Grad. Problem] Compute  $\left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)^{1+i}$  and indicate what the principal branch yields.
  - (e) Compute  $\sin(2i)$  and  $\tan(2i)$ .
  - (f) Compute  $\cosh(i\pi/3)$  and  $\sinh(i\pi/3)$ .
- #2 Exponentially Difficult Graphs Sketch each of the following figures and its image under the exponential map  $w = e^z$ . Indicate images of vertical and horizontal lines in your sketch.

Undergrads. Do (a), (c), and (e). Grads. Do them all.

- (a) The vertical strip 0 < Re(z) < 1.
- (b) The horizontal strip  $5\pi/3 < \text{Im}(z) < 8\pi/3$ .
- (c) The rectangle  $0 < x < 1, 0 < y < \pi/4$ .
- (d) [Extra Credit] The disk  $|z| \leq \pi/2$ .
- (e) [Extra Credit] The disk  $|z| \leq \pi$ .
- (f) [Extra Credit] The disk  $|z| \leq 3\pi/2$ .
- #3 Log Your Complaint Let z = x + yi be as shown below:



- (a) Is  $Log(z^2) = 2Log(z)$ ? Explain why or why not.
- (b) [Grad. Problem] In general, for which z do we have  $Log(z^2) = 2Log(z)$ ? Justify your answer.
- (c) For  $z \neq 0$ , it is always that case that  $e^{\log(z)} = z$  (and  $e^{\log(z)} = z$ ). But obviously, " $\log(e^z) = z$ " cannot hold because log is multivalued. When is it the case that  $\log(e^z) = z$ ?
- #4 A Tangential Issue One can show that  $\arctan(z) = \frac{i}{2} \log \left(\frac{1-iz}{1+iz}\right)$  when  $z \neq \pm i$ . The principal branch of inverse tangent is  $\operatorname{Arctan}(z) = \frac{i}{2} \operatorname{Log}\left(\frac{1-iz}{1+iz}\right)$  when  $z \neq \pm i$ .
  - (a) Derive the above formula for  $\arctan(z)$ .
  - (b) Show that it is never the case that  $\tan(z) = \pm i$ . [Taking into account our formula for  $\arctan(z)$ , which is defined for all  $z \neq \pm i$ , and the fact that  $\tan(\arctan(z)) = z$ , we get that the range of  $\tan(z)$  is  $\mathbb{C} \{\pm i\}$ .]