

#1 An Inverse Problem Let G be a group and let $\varphi : G \rightarrow G$ be defined by $\varphi(x) = x^{-1}$ for all $x \in G$. Show that φ is a homomorphism if and only if G is Abelian.

When G is Abelian (so that φ is a homomorphism), is φ an isomorphism? Why or why not?

Note: Since its domain and codomain match, we could call φ an endomorphism if it's a homomorphism. And we could call it automorphism if it's an isomorphism.

#2 Being Productive Let A and B be groups. Recall that $A \times B = \{(a, b) \mid a \in A \text{ and } b \in B\}$ becomes a group when we multiply coordinatewise: $(a, b)(c, d) = (ac, bd)$.

- Prove that $A \times B \cong B \times A$.
- Let $\pi_A : A \times B \rightarrow A$ be defined by $\pi_A(a, b) = a$. Show that π_A is an epimorphism (an onto homomorphism). What is its kernel? What does the first isomorphism theorem say?
- Show that $A \times B$ is Abelian if and only if *both* A and B are Abelian. Is this still true if we swap out the word “Abelian” with the word “cyclic”?

#3 Cayley Permutes Let G be a group. We write $S(G) = \{f : G \rightarrow G \mid f \text{ is a bijection}\}$ for the permutations on G . Let $L_g : G \rightarrow G$ be defined by $L_g(x) = gx$ (i.e., left multiplication by g) and $R_g : G \rightarrow G$ be defined by $R_g(x) = xg$ (i.e., right multiplication by g).

- Show that $L_a \circ R_c = R_c \circ L_a$ for all $a, c \in G$ (just plug in $b \in G$ and compute). In other words, left and right multiplication operators commute. In fact, “left and right multiplication operators commute” is equivalent to which axiom?
- Show that $L_g, R_g \in S(G)$ for every $g \in G$.
- [Grad. Problem]** Prove that $\varphi : G \rightarrow S(G)$ defined by $\varphi(g) = L_g$ is a monomorphism (i.e., one-to-one homomorphism).

Note: You showed that $L_g \in S(G)$ in the last part, so $S(G)$ makes sense as a codomain.

Observation: You just proved Cayley's Theorem: $G \cong \varphi(G)$ is a subgroup of $S(G)$ (i.e., every group is isomorphic to a subgroup of permutations).

- Consider $D_4 = \langle x, y \mid x^4 = 1, y^2 = 1, (xy)^2 = 1 \rangle = \{1, x, x^2, x^3, y, xy, x^2y, x^3y\}$. Relabel elements as integers: $1 \mapsto 1, x \mapsto 2, x^2 \mapsto 3, \dots, x^3y \mapsto 8$. Under this relabeling L_x becomes the permutation (1234)(5678). As determined by Cayley's theorem, find a subgroup of S_8 that is isomorphic to D_4 .
- [Grad. Problem]** Define $\varphi_g(x) = L_g \circ R_{g^{-1}}(x) = gxg^{-1}$ (i.e., φ_g is conjugation by g). Show φ_g is an automorphism of G .
Note: $\text{Inn}(G) = \{\varphi_g \mid g \in G\}$ is the set of *inner automorphisms* and $\text{Aut}(G)$ is the set of automorphisms. It is not hard to show that $\text{Inn}(G)$ is a subgroup of $\text{Aut}(G)$ which in turn is a subgroup of $S(G)$.
- [Grad. Problem]** Let $\varphi : G \rightarrow \text{Inn}(G)$ be defined by $\varphi(g) = \varphi_g$ (i.e., g maps to the conjugation by g map). Show that the kernel of φ is the center of G . (What does the isomorphism theorem say?) Also, show that $\text{Inn}(G)$ is a normal subgroup of $\text{Aut}(G)$.
Note: Automorphisms which aren't inner are called “outer automorphisms”. In fact, $\text{Aut}(G)/\text{Inn}(G)$ is called the *outer automorphism group* of G .

#4 Permutin' Some More Find a permutation which conjugates $\sigma = (16)(253)(4879)$ to $\tau = (149)(23)(5867)$.

#5 Quotients Write down all of the subgroups and quotients of \mathbb{Z}_{12} .

#6 Quotients Again Recall the subgroup lattice of D_4 (as given in class).

- List the normal subgroups of D_4 . For the non-normal subgroups, show why they fail to be normal by giving left coset which does not match its right coset.
- Write down all of the quotients of D_4 .